arXiv:1303.1586v2 [cond-mat.str-el] 21 May 2013

Emergent Exclusion Statistics of Fibonacci Anyons in 2D Topological Phases

Yuting Hu,[1,*] Spencer D. Stirling,[1,2,†] and Yong-Shi Wu[3,1,‡]

[1] *Department of Physics and Astronomy, University of Utah, Salt Lake City, UT 84112, USA*
[2] *Department of Mathematics, University of Utah, Salt Lake City, UT 84112, USA*
[3] *Key State Laboratory of Surface Physics, Department of Physics*
and Center for Field Theory and Particle Physics, Fudan University, Shanghai 200433, China

(Dated: May 22, 2013)

We demonstrate how the generalized Pauli exclusion principle emerges for quasiparticle excitations in 2d topological phases. As an example, we examine the Levin-Wen model with the Fibonacci data (specified in the text), and construct the number operator for fluxons living on plaquettes. By numerically counting the many-body states with fluxon number fixed, the matrix of exclusion statistics parameters is identified and is shown to depend on the spatial topology (sphere or torus) of the system. Our work reveals the structure of the (many-body) Hilbert space and some general features of thermodynamics for quasiparticle excitations in topological matter.

PACS numbers: 05.30.-d 05.30.Pr 71.10.-w 71.10.Pm

1. Introduction

By now it is well-known that (quasi-)particles in strongly entangled many-body systems may exhibit exotic quantum statistics (see, e.g., ref. 1 for a review), other than the usual Bose-Einstein and Fermi-Dirac ones. In addition to the anyonic or braiding exchange statistics[2,3] in two dimensional systems, statistical weight of many-body quantum states may also obey new combinatoric counting rules[4,5], in which the number of available single-particle states, when adding one more quasiparticle into the system, linearly depends on the number of existing quasi-particles. A typical new feature of the generalized Pauli exclusion principle is mutual exclusion between different species, resulting in a matrix of statistical parameters[4], as well as unusual thermodynamics for ideal gases with only statistical interactions[5].

More precisely, following ref. 5, in the case with only one species of quasi-particles, the number of N-particle states is assumed to be given by the binomial coefficient:

$$W_{G,N} = \binom{G_{\text{eff}} + (N-1)}{N}, \tag{1}$$

with $G_{\text{eff}} = G - \alpha(N-1)$ being the number of available single-particle states, while G is the number of single-particle states when $N = 1$. Then $\alpha = 0$ corresponds to bosons and $\alpha = 1$ fermions; other values of α gives rise to exotic exclusion statistics. Similarly, in the multi-species case, the number of many-particle states is assumed to be given by ($a, b = 1, \ldots, m$ labeling species)

$$W_{\{G_a,N_a\}} = \prod_a \binom{G_a + N_a - 1 - \sum_{b=1}^{m} \alpha_{ab}\,(N_b - \delta_{ab})}{N_a}. \tag{2}$$

Here coefficients α_{ab} form the (mutual) statistics matrix.

It has been shown[6] that the Thermodynamic Ansatz[7] for one-dimensional solvable many-particle models is actually a special case of the exotic exclusion statistics. (See also refs.8,9.) It has been also numerically verified that quasi-particle excitations in the fractional quantum Hall (FQH)systems indeed obey[10] eq. (1), or eq. (2) allowing mutual exclusion between different species[11]. Moreover either the Haldane or Jain hierarchy in the FQH effect can be theoretically understood from the exclusion statistics of quasiparticles[1,12].

Recently there has been revived interest in the study of quasiparticle statistics in 2d topological states of matter (including FQH systems), because of the possibility of using their braiding to do (fault tolerant) topological quantum computation (TQC)[13,14]. In order to know better about the error of TQC at finite temperature, it is needed to understand better how exclusion statistics of quasi-particles emerges in 2d topological matter, which governs the thermodynamics of the system.

In this paper, we carry out the many-body state counting in an exactly solvable discrete model, i.e. the Levin-Wen model[15] (with a special set of data), that describes a 2d topological quantum fluid[16] of Fibonacci anyons[17], with doubled Fibonacci anyons as fluxon excitations living on plaquettes. The Fibonacci anyons are the simplest non-abelian anyons. They occur as quasiparticles in the $k = 3$ Read-Rezayi state[18] in a FQH state with filling fraction $\nu = \frac{12}{5}$, and can be used for universal topological quantum computation[14]. (Recently, it is proposed[19] that the physics of interacting Fibonacci anyons may be studied in a Rydberg lattice gas.)

In this Letter, we first construct the number operator for fluxons in the model, which helps us identify the states with localized excitations. Then we numerically count the (many-body) states with fluxon-number N fixed, from $N = 1$ up to $N = 7$, for the system on a sphere and torus, respectively. The results exhibit a pattern closely related to the Fibonacci numbers, which in turn is put in the form of eq. (2), thus determining a topology-dependent statistical parameter matrix. Our work reveals that exotic exclusion emerges among quasiparticles due to interplay between various "hidden" degrees of freedom in addition to fluxon locations. Finally, we briefly discuss the thermodynamics of the system.

2. The model

We consider a discrete model for a "spin" system on a trivalent graph on a closed surface, e.g. a sphere or torus. We adopt a simplified formulation of the Levin-Wen model[15], with the Fibonacci data (e.g., see ref. 14) given below. Each link is assigned a "spin"-type $j = 0, 1$, and configurations of the labels on all links form an orthonormal basis in the Hilbert space. (It is conjectured that the Levin-Wen model describes a class of doubled (time-reversal invariant) topological phases[20].)

The Hamiltonian of the model is

$$\hat{H} = U\sum_v (1-\hat{Q}_v) + \epsilon \sum_p (1-\hat{B}_p), \quad \hat{B}_p = \frac{1}{D}\sum_{s=0,1} d_s \hat{B}_p^s, \tag{3}$$

where the summation runs over vertices v and plaquettes p, with $d_0 = 1$,$d_1 = \phi \equiv (\sqrt{5}+1)/2$, and $D = 1+\phi^2$. U and ϵ are positive constants.

The operator $\hat{Q}_v$ defined at vertex v is



(4)

(Only the relevant part of the graph is shown; the rest remains the same.) Here $\delta_{ijk} = \delta_{jki} = \delta_{jik}$ is given by

$$\delta_{000} = \delta_{011} = \delta_{111} = 1, \delta_{001} = 0 \tag{5}$$

(called the Fibonacci fusion rule[14]).

The operator $\hat{B}_p^s$ defined at plaquette p is



$$= v_{j_1}v_{j_2}v_{j_3}v_{j_1'}v_{j_2'}v_{j_3'}G^{j_5j_1j_3}_{sj_3'j_1'}G^{j_4j_2j_1}_{sj_1'j_2'}G^{j_6j_3j_2}_{sj_2'j_3'}, \tag{6}$$

where $v_j \equiv \sqrt{d_j}$. A similar rule applies when the plaquette p is a quadrangle, a pentagon, or a hexagon etc. Note that the matrix is nondiagonal only on the labels of the boundary links (i.e., j_1, j_2, and j_3 on the above graph).

The non-zero $6j$-symbols G are given by

$$G^{000}_{000} = 1, G^{011}_{011} = G^{011}_{111} = 1/\phi,$$
$$G^{000}_{111} = 1/\sqrt{\phi}, G^{111}_{111} = -1/\phi^2, \tag{7}$$

with the (tetrahedral) symmetry:

$$G^{ijm}_{kln} = G^{mij}_{nkl} = G^{klm}_{ijn} = G^{jim}_{lkn}. \tag{8}$$

One can check that they satisfy the conditions:

$$\text{pentagon id:} \quad \sum_{n=0,1} d_n G^{mlq}_{kpn} G^{jip}_{mns} G^{jsn}_{lkr} = G^{jip}_{qkr} G^{riq}_{mls},$$
$$\text{orthogonality:} \quad \sum_{n=0,1} d_n G^{mlq}_{kpn} G^{lmi}_{pkn} = \frac{\delta_{iq}}{d_i}\delta_{mlq}\delta_{kip}. \tag{9}$$

A notable property of the model is that $\hat{Q}_v$ and $\hat{B}_p$ are mutually commuting projection operators: $[\hat{Q}_v, \hat{B}_p] = 0$, $\hat{Q}_v\hat{Q}_{v'} = \delta_{vv'}\hat{Q}_v$ and $\hat{B}_p\hat{B}_{p'} = \delta_{pp'}\hat{B}_p$. Thus the Hamiltonian is exactly solvable. The energy eigenstates are the simultaneous eigenvectors of these projections $\hat{Q}_v$ and $\hat{B}_p$. The ground states are those $|\Phi\rangle$ that satisfy $\hat{Q}_v|\Phi\rangle = |\Phi\rangle = \hat{B}_p|\Phi\rangle$, for all v and p. Using the method developed in ref. 21, one can compute ground state degeneracy: $GSD = 1$ on sphere and $GSD = 4$ on torus.



FIG. 1: A configuration in the subspace with only fluxons allowed, with solid lines for $j = 1$ and dotted lines $j = 0$. Open strings (with only single solid line at some vertex) are forbidden by $\delta_{001} = 0$.

The quasiparticle excitations are the states with zero eigenvalue of $\hat{Q}_{v'}$ for some v' and/or of $\hat{B}_{p'}$ for some p'. In this paper we restrict ourselves to study the so-called fluxons, satisfying $\hat{Q}_v = 1$ for all v, $\hat{B}_{p'} = 0$ for a specified set of p' (where fluxons live), and $\hat{B}_p = 1$ for all other p. (See Fig. 1.)

3. Number operator of fluxons

A crucial property of the model is that $\hat{B}_p^s$ defined above form an abelian algebra:

$$\hat{B}_p^r \hat{B}_p^s = \sum_{t=0,1} \delta_{rst} \hat{B}_p^t, \tag{10}$$

which can be verified using conditions (8) and (9). From the generators we construct the operators ($i = 0, 1$):

$$\hat{n}_p^i = \sum_{j=0,1} S_{i0} S_{ij} \hat{B}_p^j, \tag{11}$$

where S is the modular matrix given by

$$S = \frac{1}{\sqrt{D}} \begin{pmatrix} 1 & \phi \\ \phi & -1 \end{pmatrix}, \tag{12}$$

with S_{ij} for fixed i being a one-dimensional representation of the algebra (10). One can check that $\hat{n}_p^i$ ($i = 0, 1$) form a complete set of orthonormal projections:

$$\hat{n}_p^i \hat{n}_p^k = \delta_{ik} \hat{n}_p^k, \quad \hat{n}_p^0 + \hat{n}_p^1 = \mathbf{1}. \tag{13}$$

There is a fluxon at p in a state $|\Psi\rangle$, if $\hat{n}_p^1|\Psi\rangle = |\Psi\rangle$. A ground state $|\Phi\rangle$ contains no fluxon because $\hat{n}_p^0 = \hat{B}_p$. Hence the model has only one type of fluxons. No state contains two fluxons at any p. However, we will show that the fluxon in this model is *not* a fermion; rather its state counting needs exotic exclusion statistics.

4. Exclusion Statistics On a sphere

Let us count the N-fluxon states in the model with P plaquettes on a sphere. Pick up a set of N fixed plaquettes and denote it by $\mathcal{C} = \{p_1, p_2, ..., p_N\}$ ($N < P$).

The states with exactly N fluxons occupying the selected plaquettes are those $|\psi\rangle$ satisfying

$$\begin{aligned} \hat{n}_p^j|\Psi\rangle &= \delta_{j1}|\Psi\rangle, \quad \text{for } p \in \mathcal{C}, \\ \hat{n}_{p'}^j|\Psi\rangle &= \delta_{j0}|\Psi\rangle, \quad \text{for } p' \notin \mathcal{C}. \end{aligned} \tag{14}$$

Thus $\left(\prod_{p\in\mathcal{C}} \hat{n}_p^1 \prod_{p'\notin\mathcal{C}} \hat{n}_{p'}^0\right)$ is the projector onto the subspace of such states. Tracing this projection computes the total number of the N-fluxon states in the configuration $\mathcal{C}$:

$$w_{P,N,\mathcal{C}} = \mathrm{tr}(\prod_{p\in\mathcal{C}} \hat{n}_p^1 \prod_{p'\notin\mathcal{C}} \hat{n}_{p'}^0). \tag{15}$$

We numerically compute eq. (15) on random graphs on sphere with $P(\geq 7)$ plaquettes, with the stable result presented in Table I.

TABLE I: State Counting on Sphere

Fluxon number N	0	1	2	3	4	5	6	7
State Counting $w_{P,N,\mathcal{C}}$	1	0	1	1	4	9	25	64

The pattern of the N-dependence is obvious:

$$w_{P,N,\mathcal{C}} = F_{N-1}^2, \tag{16}$$

where F_n is the Fibonacci number that satisfies the recurrence relation: $F_n = F_{n-1} + F_{n-2}$ with $F_1 = F_2 = 1$. Both numerically and analytically we have checked that eq. (16) is independent of the graph, of the total number P of plaquettes, as well as the locations of the N fluxons. The appearance of the squared in eq. (16) is consistent with the conjecture that the LW model describes a *doubled* topological phases[21].

Summing over configurations $\mathcal{C}$ (i.e. over possible distributions of N plaquettes in a fixed graph), we get the total number of N-fluxon states:

$$W_{P,N}^{\text{sphere}} = \sum_{\mathcal{C}} w_{P,N,\mathcal{C}} = \binom{P}{N} F_{N-1}^2. \tag{17}$$

The first factor counts the ways to distribute N fluxons over P plaquettes. The second factor counts the configurations of the link degrees of freedom, which are not unique, given N and $\mathcal{C}$. The independence of $w_{P,N,\mathcal{C}}$ on P and $\mathcal{C}$ implies the degeneracy of the excited states is topological in the sense that it does not depend on the detailed structure of the underlying graph, and not on the relative positions between the fluxons as well. The origin of this property lies in the topological symmetry of the model under mutations of the underlying graph[21].

To find the exclusion statistics, we rewrite (17):

$$W_{P,N}^{\text{sphere}} = \binom{P}{N} \sum_{N_1,N_2=0}^{[\frac{1}{2}(N-2)]} \binom{N-N_1-2}{N_1} \binom{N-N_2-2}{N_2}, \tag{18}$$

where $[x]$ is the greatest integer less than or equal to x.

Now eq. (2) is of the form of eq. (18), by introducing two additional pseudo-species $a = 1, 2$, which do not contribute to the total energy but are helpful for state-counting. This is similar to what was suggested for state counting in some conformal field theories[22]. Including the original fluxon species labeled by $a = 0$, from eq. (18) we read the exclusion statistics parameters α_{ab} $(a, b, = 0, 1, 2)$:

$$\alpha^{\text{sphere}} = \begin{pmatrix} 1 & 0 & 0 \\ -1 & 2 & 0 \\ -1 & 0 & 2 \end{pmatrix}. \tag{19}$$

The diagonal α_{aa} is the self-exclusion statistics for species a. The $\alpha_{00} = 1$ implies the hard-core boson behavior. This can be understood with eq. (13).

The pseudo-species provides a way to count configurations, in the presence of fluxons, of link degrees of freedom, which are not uniquely determined by the constraints (14). The value $\alpha_{11} = \alpha_{22} = 2$ implies that one pseudo-particle makes two single-particle states (or "seats") unavailable to an additional pseudo-particle. The negative mutual statistics $\alpha_{20} = \alpha_{30} = -1$ tells us that each fluxon present creates one vacant "seat" for each pseudo-species. So the maximum particle number of each pseudo-species is naturally $[(N-1)/2]$. These results help us to understand the structure of the (many-body) Hilbert space for excited states of the system, and to derive analytically the state counting formula (18).

6. Exclusion statistics on a torus

We proceed to consider the model on a torus. The ground state degeneracy[21] is 4 . Thus the system exhibits the global topological degrees of freedom, and we can study their effects on excited states by counting the pseudo-particle states.

Pick up N plaquettes ($N < P$). The number of states with N fluxons on htese plaquettes is computed numerically as in Table II.

TABLE II: State Counting on Torus

Fluxon number N	0	1	2	3	4	5	6
State Counting	2^2	1	3^2	4^2	7^2	11^2	18^2

The pattern of its dependence on N is

$$W_{P,N}^{\text{torus}} = \binom{P}{N} L_N^2, \tag{20}$$

with L_n the Lucas number, a modified version of the Fibonacci number, satisfying the recurrence relation $L_n = L_{n-1} + L_{n-2}$ with $L_1 = 1, L_2 = 3$.

We rewrite (20) in terms of binomial coefficients:

$$W^{\text{torus}}_{P,N} = \binom{P}{N} \sum_{N_1,N_2=0,1} \binom{1}{N_1}\binom{1}{N_2} \times \sum_{N_3,N_4=0}^{[\frac{1}{2}(N-2)]} \binom{N-2N_1-N_3}{N_3}\binom{N-2N_2-N_4}{N_4}, \tag{21}$$

and get the exclusion statistics parameters α_{ab} $(a, b = 0, 1, 2, 3, 4)$:

$$\alpha^{\text{torus}} = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ -1 & 2 & 0 & 2 & 0 \\ -1 & 0 & 2 & 0 & 2 \end{pmatrix}, \tag{22}$$

where we denote by $a = 0$ the fluxon species.

Eq. (21) shows that one needs to introduce four pseudo-species $a = 1, 2, 3, 4$. The pseudo-species $a = 1, 2$ are interpreted as the topological degrees of freedom on the torus, for the following reasons. The allowed "particle number" $N_1, N_2 = 0, 1$ of these pseudo-species are independent of the number N of fluxons. Particularly when there is no fluxon present, the configurations $N_1, N_2 = 0, 1$ characterize the four-degenerate ground states. Then the pseudo-species $a = 3, 4$ provide a way to count the configurations of link degrees of freedom given a ground state and fluxon number.

The state counting of excitations on a torus is shown different from that on a sphere. Indeed the mutual statistics parameters $\alpha_{31} = \alpha_{42} = 2$ imply that the number of configurations of link degrees of freedom $a = 3$ $(a = 4)$ are affected by the topological degrees of freedom $a = 1$ $(a = 2)$, respectively. On the other hand, the topological degrees of freedom are not affected by the fluxons present and the link degrees of freedom. So the degenerate ground states can be used to label the sectors of excitations. We note that in the sector with $N_1 = N_2 = 1$, the state counting for fluxons is exactly the same as that on sphere.

7. Statistical Thermodynamics

Now we assume that only fluxons can be thermally excited; this is the case when $U >> \epsilon, kT$ in eq. (3). In the thermodynamic limit, the Hilbert space dimension of N-fluxon states (occupying N fixed plaquettes) is asymptotically

$$\begin{aligned} &\text{on sphere:} \qquad \lim_{N\to\infty} F^2_{N-1} \sim \phi^{2N-2}/5, \\ &\text{on torus:} \qquad \lim_{N\to\infty} L^2_{N} \sim \phi^{2N}. \end{aligned} \tag{23}$$

(ϕ^2 is called the quantum dimension of the fluxon.) On a torus, for example, the canonical partition function is

$$Z^{\text{torus}} = \sum_{N=0}^{P} \binom{P}{N} L^2_N e^{-N\epsilon/kT} \sim (\phi^2 e^{-\epsilon/kT} + 1)^P. \tag{24}$$

It can be interpreted as the grand canonical partition function of the many-fluxon system, which behaves like a fermionic system with a *temperature-independent fugacity* z given by the quantum dimension:

$$z = \phi^2. \tag{25}$$

The fugacity z counts the effective number of states per fluxon located at a plaquette. Note that z is irrational rather than integer. This is a manifestation that the many-fluxon states are highly entangled ones with long-range entanglement. They are superpositions of highly constrained j-configurations on the links, obviously not of the form of a direct product of localized fluxon states.

The statistical distribution of the average occupation number of fluxons is obtained from eq. (24):

$$\langle n \rangle = \langle N \rangle / P = \frac{1}{e^{\epsilon/kT}\phi^{-2} + 1}. \tag{26}$$

Many useful thermodynamic observables are then computable. Though the model is very simple, we believe that the features revealed in this paper should be quite general for emergent exotic exclusion statistics and thermodynamics for quasiparticle excitations in a wide class of 2d topological phases.

Acknowledgement: YH thanks Department of Physics, Fudan University for warm hospitality he received during a visit in summer 2011 and 2012. YSW was supported in part by US NSF through grant No. PHY-1068558.

Note added. After this paper was posted on arXiv, we learn that a different state-counting formula, without proof and connection to exclusion statistics, appeared in M. D. Schulz, S. Dusuel, K. P. Schmidt, and J. Vidal, arXiv:1212.4109.

---

[*] Electronic address: yuting@physics.utah.edu

[†] Electronic address: stirling@physics.utah.edu

[‡] Electronic address: wu@physics.utah.edu

[1] Y.S. Wu, "*Fractional Statistics and Strongly Correlated Systems*", in "*Topics in Theoretical Physics*", ed. Y.M. Cho, (World Scietific, 1996); pp. 27-59.

[2] F. Wilczeck, Phys. Rev. Lett. **48**, 1144 (1982).

[3] Y.S. Wu, Phys. Rev. Lett. **52**, 2103 (1984).

[4] F.D.M. Haldane, Phys. Rev. Lett. **67**, 937 (1991).

[5] Y.S. Wu, Phys. Rev. Lett. 73, 922 (1994).

[6] D. Bernard, Y.S. Wu, in "*New Developments of Integrable Systems and Long-Range Interaction Models*", ed. M.L Ge

and Y.S. Wu, (World Scientific, 1995); pp. 10-21.

[7] C.N. Yang, C.P. Yang, J. Math. Phys. **10**, 1315 (1969).

[8] Z.N.C. Ha, Nucl. Phys. **B435** 604 (1995).

[9] Y. Hatsugai, M. Kohmoto, T. Koma, Y.S. Wu, Phys. Rev. **B 54**, 5358 (1996).

[10] S. He, X.C. Xie, F.C. Zhang, Phys. Rev. Lett. **68**, 3460 (1992).

[11] W.P. Su, Y.S. Wu, J. Yang, Phys. Rev. Lett. **77**, 3423 (1996).

[12] Y.S. Wu, Y. Yu, Y. Hatsugai, and M. Kohmoto, Phys. Rev. **B 57**, 9907 (1998).

[13] A. Yu. Kitaev, Ann. Phys. **303**, 2 (2003).

[14] Z. Wang, "*Topological Quantum Computation*", CMBS no. 112, American Mathematical Society (Providence, RI; 2010).

[15] M. Levin, X.G. Wen, Phys. Rev. **B 71**, 045110 (2005).

[16] C. Gils, S. Trebst, A. Kitaev, A. Ludwig, M. Troyer, Z.H. Wang, Nature Physics **5**, 834 (2009).

[17] J.K. Slingerland, F.A. Bais, Nucl. Phys. **B 612**, 229 (2001).

[18] N. Read, E. Rezayi, Phys. Rev. **B 59**, 8084 (1999).

[19] I. Lesanovsky, H. Katsura, Phys. Rev. **A 86**, 041601(R) (2012).

[20] M. Freedman, C. Nayak, K. Shtengel, K. Walker, Z. Wang, Ann. Phys. **310**, 428 (2004).

[21] Y. Hu, S. Stirling, Y.S. Wu, Phys. Rev. **B 85**, 075107 (2012).

[22] S. Guruswamy, K. Schoutens, Nucl. Phys. **B 556**, 530 (1999).